



DEB-in-Practice


DEB tools to Trait Based

Ecology

Starrlight Augustine
starrlight@tecnico.ulisboa.pt
University of Lisboa/ Vrije
Universiteit

Dina Lika
lika@uoc@gr
University of Crete

Inter-species comparisons



- parameter values
- implied properties

Know your labels !!!

Careful with temperature !!!



Example to be found: GitHub [add-my-pet](#) / [SI](#)

AmPtool


AmP data

Manual: amptool.debtheory.org/docs/

Five taxonomic groups:




What traits co-vary?



- Primary scaling relationships
 - Covariation of primary parameters
- Secondary scaling relationships
 - Covariation of compound parameters
- Tertiary scaling relationships
 - Deal with phenomena at larger scales in time and space


Primary scaling relationship



Plot maturity level at birth against maximum structural


Each of the 5 groups works with different taxa (10 min)

Then come to board and present the different graphs




Mammalia; Crustacea; Chondrichthyes; Actinopterygii; Reptilia

Primary scaling: coding support:



1st define your legend; order matters

```
legend_mamm = { ... %
    {'o', 8, 1, [0 0 0], [0 1 1]}, 'Mammalia'
    {'o', 8, 1, [1 1 1], [0.8 0.8 0.8]}, 'Animalia'
};
```



The order matters: Animalia in the back, and Mammalia in the front


```
>> lineage('Mammalia')
```

```
ans =
```

```
16x1 cell array
```

```
{'Animalia'      }
{'Eumetazoa'     }
{'Bilateria'     }
{'Nephrozoa'      }
{'Deuterostomata' }
{'Chordata'       }
{'Olfactores'     }
{'Vertebrata'     }
{'Gnathostomata'  }
{'Osteichthyes'   }
{'Sarcopterygii'  }
{'Rhipidistia'    }
{'Tetrapoda'      }
{'Amniota'        }
{'Synapsida'      }
{'Mammalia'       }
```


amptool.debtheory.org/docs/



```
shstat_options('default');
LiEHb = read_allStat({'L_i', 'E_Hb'}); 
```

```
[Li_EHb, leg] = shstat(LiEHb, legend_mamm, 'Mammalia');
figure(Li_EHb)
xlabel('_{10}log ultimate struc length, L_i^\infty, cm')
ylabel('_{10}log E_H^b, J')
print -r300 -dpng Li_EHb_mammalia.png
figure(leg)
print -r300 -dpng leg_mammalia.png
```


Primary scaling relationship



Plot maturity level at birth against maximum structural



Each of the 5 groups works with different taxa (10 min)

Then come to board and present the different graphs




Mammalia; Crustacea; Chondrichthyes; Actinopterygii; Reptilia


Primary scaling





Primary scaling



Chondrichthyes vs Actinopterygii




Did you know?




```
>> lineage('Sarcophilus_harrisii')
```

```
{'Animalia'}  
{'Eumetazoa'}  
{'Bilateria'}  
{'Nephrozoa'}  
{'Deuterostomata'}  
{'Chordata'}  
{'Olfactores'}  
{'Vertebrata'}  
{'Gnathostomata'}  
{'Osteichthyes'}  
{'Sarcopterygii'}  
{'Rhipidistia'}  
{'Tetrapoda'}  
{'Amniota'}  
{'Synapsida'}  
{'Mammalia'}  
{'Theriiformes'}  
[{'Marsupialia'}]  
{'Dasyuromorphia'}  
{'Dasyuridae'}  
{'Dasyurinae'}  
{'Dasyurini'}  
{'Sarcophilus'}  
{'Sarcophilus_harrisii'}
```



amptool.debtheory.org/docs/



```
treeview_taxa('Marsupialia')
```

```
prtStat('Marsupialia', 'p_M');
```

```
pedigree('Marsupialia');
```


```

legend_mamm = { ... %


{'o', 8, 1, [0 0 0], [1 1 0]}, 'Marsupialia'
{'o', 8, 1, [0 0 0], [0 1 1]}, 'Mammalia'
{'o', 8, 1, [1 1 1], [0.8 0.8 0.8]}, 'Animalia'

};



```



add a level to the legend and then rerun the code for the plot and the extra group appears !!!




Secondary scaling relationship



$$[E_m] = \frac{\{P_{Am}\}}{\nu} = z[E_m^{\text{ref}}]$$


A large-bodied species has larger reserve capacity

Inter-species comparison In practice:



Plot maximum reserve density again maximum length - secondary

Each of the 5 groups works with different taxa (10 min)
Then come to board and present the different graphs









```
shstat_options('default');
LiEm = read_allStat('L_i', 'E_m');
```




```
[Li_Em, leg] = shstat(LiEm, legend_mamm, 'Mammalia');
figure(Li_Em)
xlabel('_{10}log ultimate struc length, L_i^{\infty}, cm')
ylabel('_{10}log maximum reserve density, [E_m], J/cm^3')
print -r300 -dpng Li_Em_mammalia.png
figure(leg)
print -r300 -dpng leg_mammalia.png
```

Secondary scaling




Secondary scaling relationship



von Bertalanffy growth rate:


$$r_B = \frac{\dot{k}_M / 3}{1 + f / g} = \frac{\dot{k}_M^{ref} / 3}{1 + zf / g_{ref}}$$



$$V^{1/3}(a) = V_\infty^{1/3} - (V_\infty^{1/3} - V_b^{1/3}) \exp(-r_B a)$$


\dot{r}_B	von Bertalanffy growth rate
\dot{k}_M	somatic maintenance rate coeff
g	energy investment ratio
z	zoom factor

Inter-species comparison In practice:



Plot von Bert as function of maximum length with and without temperature correction

10 minutes





```
shstat_options('default');
LirBcT = read_allStat({'L_i', 'r_B', 'c_T'});
    LirB = [LirBcT(:,1), LirBcT(:,2)];
[Li_rB, leg] = shstat(LirB, legend_mamm, 'Mammalia');
figure(Li_rB)
xlabel('_{10}log ultimate struc length, L_i^\infty, cm')
ylabel('_{10}log r_B, 1/d, T_typical')
print -r300 -dpng Li_rB_mammalia.png
figure(leg)
```




Very important: These quantities like `r_B` and `E_m` are read from AmPdata structure !!! Food dependant statistics are all given at `T_typical`

Secondary scaling temperature correction - values at T_typical



amptool.debtheory.org/docs/





Create same plot but put the VB growth rate at reference temperature for all species:

```
shstat_options('default');
LirBcT = read_allStat({'L_i', 'r_B', 'c_T'});
LirB = [LirBcT(:,1), LirBcT(:,2)./ LirBcT(:,3)];
[Li_rB, leg] = shstat(LirB, legend_mamm, 'Mammalia');
figure(Li_rB)
xlabel('_{10}log ultimate struc length, L_i^\infty, cm')
ylabel('_{10}log r_B, 1/d, T_ref')
print -r300 -dpng Li_rB_mammalia_Tref.png
figure(leg)
```





Very important: These quantities like r_B and E_m are read from AmPdata structure !!! Food dependant statistics are all given at $T_{typical}$


Secondary scaling temperature correction - values at T_reference



Secondary scaling- von Bert growth rate- T_{ref} - by taxa




Inter-species comparison In practice:



- For example: weight at birth as function of ultimate weight.
- You can calculate quantities you do not have data for to check consistency.
- You have to think about the levels of organisation in the phylogeny to make meaningful plots
- Be careful with temperature!!!

Inter-species comparison In practice:




debportal.debtheory.org/docs/DEBpapers.html#Patterns

There is a link to the add-my-pet/SI GitHub repository where the matlab code that generate figures for all of the papers are provided

You can use this for inspiration

Survivor or survival function




Let T be a continuous random variable with probability density function $f(t)$ and cumulative distribution function $F(t)$ on the interval $[0, \infty)$.


The survivor function $S(t)$ is

$$S(t) = P(T > t) = \int_t^{\infty} f(u)du = 1 - F(t)$$

It gives the probability that an object will survive past a certain time or the proportion of the values of T that are larger than t

The probability of taking values larger than 4 is 0.4;
 $S(4) = 0.4$






Empirical survivor function

The empirical survivor function is the distribution function associated with the values of a sample. It is a step function that jumps down by $1/n$ at each of the n data points. Its value at any specified value of the measured variable is the fraction of observations of the measured variable that are greater than or equal to the specified value.



DEDtool function `surv` computes the survival function

Example

```
xy = surv([3 8 2 1 4 5 6 9]);  
plot(xy(:,1), xy(:,2), 'b')
```




Empirical survivor function




80% of Aves have $\kappa > 0.95$

80% of Actinopterygii have $\kappa > 0.75$


95% of Aves and 67% of Actinopterygii have κ values larger than 0.8



Survival function of compound parameter



maximum reserve density




Survival function coding support:

% line legends:







```
llegend_mamm = { ... %  
  { '-' , 2, [0 1 1] } , 'Mammalia'  
  { '-' , 2, [0.8 0.8 0.8] } , 'Animalia'  
};  
↑  
line style  
↑  
line width  
↑  
color
```

Survival function coding support:




```
shstat_options('x_transform', 'log10');
hE_m = shstat({'E_m'}, llegend_mamm, 'Mammalia');
figure(hE_m)
xlabel('_{10}log reserve capacity, [E_m], J/cm^3')
print -r300 -dpng Em_mamm.png
```

Distribution of maximum reserve density



You can also query by ecocodes:



```
select_eco('ecozone', { 'MS', 'TS' })
```

Selection of all (terrestrial and marine) Antarctic species

```
speciesNames = select_eco('ecozone', { 'MS', 'TS' });
```

```
LiPM_antartica = read_stat(speciesNames, { 'L_i', 'p_M' });
plot(log10(LiPM_antartica(:,1)), log10(LiPM_antartica(:,2)), 'ro')
```

Just to show that you can make your own plots.... many plots possible but they do not always make sense. Notice that the temperature is at **T reference**. Taken from AmPdata !

Thank you



Questions?